

Is QGP the free quark/gluon gas ? – Lattice QCD Study of meson correlators near the phase transition –

Ph. de Forcrand^a M. García Pérez^b T. Hashimoto^c S. Hioki^d
H. Matsufuru^e O. Miyamura^f A. Nakamura^g
I.-O. Stamatescu^{h,i} T. Takaishi^j T. Umeda^f

^a*ETH-Zürich, CH-8092 Zürich, Switzerland*

^b*Dept. Física Teórica, Universidad Autónoma de Madrid, E-28049 Madrid, Spain*

^c*Department of Applied Physics, Faculty of Engineering Fukui University,
910-8507, Japan*

^d*Department of Physics, Tezukayama University, Nara 631-8501, Japan*

^e*Research Center for Nuclear Physics, Osaka University, Ibaraki 567-0047, Japan*

^f*Department of Physics, Hiroshima University, Higashi-Hiroshima 739-8526,
Japan*

^g*RIISE, Hiroshima University, Higashi-Hiroshima 739-8521, Japan*

^h*Institut für Theoretische Physik, Universität Heidelberg, D-69120 Heidelberg,
Germany*

ⁱ*FEST, Schmeilweg 5, D-69118 Heidelberg, Germany*

^j*Hiroshima University of Economics, Hiroshima 731-01, Japan*

Presented by: A. Nakamura

Abstract

We report the study of hadron properties near the confinement/deconfinement transition temperature T_c obtained by quenched lattice QCD simulations. We analyze *temporal* and *spatial* meson correlators and the wave functions. Below T_c , we observe little change in the meson properties as compared with $T = 0$. Above T_c we observe new features: chiral symmetry restoration and signals of plasma formation, but also indication of persisting “mesonic” (metastable) states and different temporal and spatial “masses” in the mesonic channels. This suggests a complex picture of QGP in the region near the phase transition.
